

Phase Control of Laser-Matter Interactions and Applications

Prof. Marcos Dantus

Prof. Vadim V. Lozovoy

Prof. Gavin Reid

Prof. Dan Jones

Prof. Frank Wise

Yves Coello

Bai Nie

Ilyas Saytashev

Anton Ryabtsev

Christine Kalcic,

Jay Shah

Arkaprabah Konar

Marshall Bremer

Xin Zhu

Jess Gun





Paul Wrzesinski

Johanna De la Cruz

Janelle Shane

Nelson Winkler

Matthew Comstock



Biophotonic
SOLUTIONS INC.
ULTRAFAST PULSE SHAPING & COMPRESSION

Dr. Igor Pastirk

Dr. Tissa Gunaratne

Dr. Bingwei Xu

Dr. Dmitry Pestov



COI Disclosure: Dantus is the founder and president of Biophotonic Solutions Inc.

2001

A 6fs pulse Bardeen and Shank in 1987; Sub-6fs Baltuska and Weirsmas 1997

Computer-controlled phase amplitude pulse shapers demonstrated 1994,
Commercially available after 2001.

NEW VOICES IN CHEMISTRY

MARCOS DANTUS

LASER CONTROL OF CHEMICAL REACTIONS

Scientific and technological breakthroughs
breathe renewed life into area of study



*“Smart photons, from ultrafast shaped pulses, could revolutionize techniques such as **laser machining**, laser deposition, and laser desorption.”*

*“The ultimate goal of this research is to produce an **automated protein-sequencing apparatus**.”*

Also accomplished in early 2000's

- Automated pulse compression
- Coherent communications with fs pulses
- Functional imaging to detect cancer
- Detection of warfare agents

M. Dantus, Chemical & Engineering News **79**, 191-191 (2001)

Outline:

1. Introduction

2. Progress in Quantum Control


- a. Wave packet manipulation
- b. Interference between laser pulses and wave packets
- c. Selective two-photon excitation, CARS, SRS, electronic and vibrational control

3. Challenges in Chemistry

- a. Chemical reactions
- b. Coherent Control of Electrons, Electronic States and Electronic Transitions
- c. Breaking strong chemical bonds while leaving weak ones intact
- d. Determining the role of coherent quantum processes in nature

4. Future Outlook and Applications

- a. Phase Control of Advanced Sources
- b. Quantum control-based molecular sensing and diagnostics
- c. Material Synthesis and Material Processing
- d. Connecting the Macro-Micro-and-Nano Worlds
- e. Coherent control and communications and information processing. Internet is incoherent



A Brief History of Quantum Phase Control

- Young's double-slit experiment (1803), although it is just linear interference

$$P \propto \left| e^{i\phi_1} + e^{i\phi_2} \right|^2 \propto \cos(\phi_1 - \phi_2)$$

- Brumer and Shapiro propose one vs. three-photon quantum control (1986)

$$P \propto \left| e^{i\phi_3} + e^{i3\phi_1} \right|^2 \propto \cos(\phi_3 - 3\phi_1) \text{ (One final } 3\omega_0 \text{ narrow frequency)}$$

- Weiner demonstrates Optical Code Division Multiple Access (CDMA) (1988, 1990, 1998)
(using inversion symmetry of binary phase codes to maximize SHG at $2\omega_0$)

- Gordon Experimental demonstration of one vs. three-photon control (1997)

- Silberberg demonstrates multi-chromatic two-photon control (1998)

$$P(\Delta) \propto \left| \int e^{i[\phi(\omega-\Delta)-\phi(\omega+\Delta)]} d\omega \right|^2 \text{ (One final } \Delta \text{ narrow frequency)}$$

$$P(2\omega_0) = \left| \int e^{i[\phi(\omega_0-\Delta)+\phi(\omega_0+\Delta)]} d\Delta \right|^2 \text{ (One final narrow frequency } 2\omega_0)$$

- Weinacht, Ahn and Bucksbaum “write-read” on sculpted Rydberg wave packet (1998)

MII for other nonlinear processes


Two-photon excitation

$$E^{(2)}(2\omega) \propto \int_{-\infty}^{\infty} |E(\omega + \Omega)| |E(\omega - \Omega)| \exp\left[i\{\varphi(\omega + \Omega) + \varphi(\omega - \Omega)\}\right] d\Omega.$$

N-photon excitation




$$E^{(n)}(n\omega) \propto \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} d\Omega_1 \dots d\Omega_{n-1} |E(\omega + \Omega_1)| \dots |E(\omega + \Omega_{n-1})| |E(\omega - \Omega_1)| \dots |E(\omega - \Omega_{n-1})| \times \\ \exp\left\{i\left[\varphi(\omega + \Omega_1) + \dots + \varphi(\omega + \Omega_{n-1}) + \varphi(\omega - \Omega_1) + \dots + \varphi(\omega - \Omega_{n-1})\right]\right\}$$

Selective two-photon excitation, SHG or SRS using Binary Phases


$$E^{(2)}(2\omega) \propto \int_{-\infty}^{\infty} E(\omega + \Omega) E(\omega - \Omega) d\Omega = \int_{-\infty}^{\infty} \exp\left[i\{\varphi(\omega + \Omega)\}\right] \exp\left[i\{\varphi(\omega - \Omega)\}\right] d\Omega = \sum E_i E_j$$


J.Phys. Chem. A, 106, 9369, 2002. J.Chem. Phys. **118**, 3187 (2003) Opt. Express, 12, 1061 (2004).
ChemPhysChem, 6, 1952 (2005), Physical Review A **74**, 041805(R) (2006)

“An atomic switch!” (P. Bucksbaum, Nature)



Two-photon excitation of Cs atoms vs. position of a step phase function




Two-photon excitation vs. position of a step phase function. No enhancement observed at 0 detuning.

D. Meshulach, Y. Silberberg, *Nature* 1998, 396, 239-242

Lozovoy and Dantus multichromatic multifrequency n-photon control (2002-)

Multiphoton Intrapulse Interference


$$I_{SHG}(2\omega) \propto \left| \int |E_0(\omega + \Omega)| |E_0(\omega - \Omega)| \exp[i(\varphi(\omega + \Omega) + \varphi(\omega - \Omega))] d\Omega \right|^2$$



J.Phys. Chem. A, 106, 9369, 2002. J.Chem. Phys. **118**, 3187 (2003) Opt. Express, 12, 1061 (2004).
ChemPhysChem, 6, 1952 (2005), Physical Review A **74**, 041805(R) (2006)

Brief Review of Pulse Shaping Technology

Time Domain




Directly shape $E(t)$
using ultrafast processes

Stavola, Agostinelli, Sceats, *Appl. Opt.* **18**, 4101 (1979)

J. G. Fujimoto and E. P. Ippen, *Opt. Lett.* **8**, 446 (1983)

Bartels, Weinacht, Wagner, Baertschy, Green, Murnane, Kapteyn, *Phys. Rev. Lett.* **88**, 013903 (2002)

Frequency Domain



$$E(t) \xrightarrow{\text{Fourier Transform}} E(\omega)$$

Shape

$$E_{out}(\omega) \xrightarrow{\text{Inverse Transform}} E_{out}(t)$$

Desbois, Gires, Tourniois, *IEEE J. Quantum Electron.* **QE-9**, 213 (1973)

Agostinelli, Harvey, Stone, Gabel, *Appl. Opt.* **18**, 2500 (1979)

Froehly, *J. Opt. (Paris)*, **12**, 25 (1981)
"The Fourier Plane"

Weiner and Heritage, *Rev. Phys. Appl.* **22**, 1619 (1987)


Weiner, Leaird, Patel, Wullert, *Opt. Lett.* **15**, 326 (1990)

Wefers, Nelson *Opt. Lett.* **20**, 1047 (1995)

Commercially available:

Biophotonic Solutions Inc.

Freq. and Time Domain



Shape a combination of
 $E(t)$ and $E(\omega)$

Haner and Warren, *Appl. Phys. Lett.* **52**, 1458 (1988)


Fermann, da Silva, Smith, Silberberg, Weiner, *Opt. Lett.* **18**, 1505 (1993)

Verlueise, Laude, Cheng, Spielmann, Tourniois, *Opt. Lett.* **25**, 575 (2000)

Commercially available:

Fastlite

Multiphoton Intrapulse Interference Phase Scan (MIIPS)



Opt. Lett. 29, 775 (2004)

JOSA B 23, 750 (2006)



Laser Focus World 43, 101, (2007)

JOSA B 25, A140 (2008)


MIIPS based techniques and their applications are protected by 8 issued and 25 patents pending.

Multiphoton Intrapulse Interference Phase Scan (MIIPS)

GVD Measurements With $\pm 0.1 \text{ fs}^2$ Accuracy and Precision



Dispersion of water



Coello *et al.*, *Applied Optics* **46**, 8394 (2007), Water and Sea Water

Devi, *et al.* *AIP Advances* **1**, 032166 (2011), Common Solvents


Wrzesinski, *et al.* *Opt. Express* **19**, 5163 (2011), Atmospheric and Combustion Gases

2. Progress in Quantum Control

a. Wave Packet Manipulation and Role of Homogeneous and Inhomogeneous Broadening


1-photon pump 2-photon probe

Inner vs. outer turning point probing to different states of I_2



Rice, Tannor, Kosloff
Dantus, Zewail, Gerber,
1985-88


Time domain approach,
intuitive results.



Bowman Dantus Zewail, Chem. Phys. Lett. 174, 546 (1990)

2-photon pump 1-photon probe

Inner vs. outer turning point probing to different states of Na_2




Na^+ vs Na_2^+ Baumert, Gerber, Isr.J.Chem. 34, 103 (1994)




2. Progress in Quantum Control

a. Control of ground and excited state coherence cancelling inhomogeneous broadening at room temp.

a) Homogeneous vibronic relaxation



b) Inhomogeneous vibronic relaxation




Pastirk et al. Chem.Phys.Letts. **333**, 76 (2001)


Brown et al. J. Chem. Phys. **111**, 3779 (1999)

Dantus, Annu. Rev. Phys. Chem. **52**, 639 (2001)

Lozovoy Dantus Chem.Phys.Letts. **351**, 213 (2002)



Intra- and intermolecular coherence, the role of local fields






Lozovoy et al, J. Chem. Phys. **267**, 99 (2001)


2. Progress in Quantum Control

b. Interference Between Laser Pulses and Wave Packets

Scherer et al. JCP 93, 856 (1990), JCP 95, 1487 (1991)




Michelson Interferometer



1st order autocorrelation



2nd order autocorrelation




2. Progress in Quantum Control

b. Interference Between Laser Pulses and Wave Packets


Multiple independent comb shaping (MICS)
Avoiding linear interference by pulse shaping

Theory



MICS: Pestov et al. *Opt. Exp.* 17, 14351 (2009)
IR144: Konar et al. *J. Phys. Chem.* (2012)


Experimental
IR144 in solution




Fluorescence



Stim. Emission



Stim. Emission





Fluorescence


2. Progress in Quantum Control

c. Selective Two-Photon Excitation, CARS, SRS, Electronic and Vibrational Control


Single-pulse CARS with selective vibrational mode excitation: Dudovich Oron Silberberg Nature 418, 512 (2002)



Sinusoidal modulation




Binary Phase Modulation: Physical Review A **74**, 041805(R) (2006)



2. Progress in Quantum Control

c. Selective Two-Photon Excitation, CARS, SRS, Electronic and Vibrational Control




J. Phys. Chem. 108, 53 2004



Selective two-photon Microscopy
Optics Express 11, 1695 (2003)

Live fluorescently labeled
Drosophila embryos


Jennifer P. Ogilvie et al.
Optics Express 14, (2006)



Highly specific label-free molecular imaging with spectrally tailored excitation-stimulated Raman scattering (STE-SRS) microscopy




Christian W. Freudiger^{1,2}, Wei Min², Gary R. Holtom², Bingwei Xu³, Marcos Dantus⁴
and X. Sunney Xie^{2*}

Nature Photonics, 294 (2010)



2. Progress in Quantum Control

c. Selective Two-Photon Excitation, CARS, SRS, Electronic and Vibrational Control





Dela Cruz et al. *Opt. Express*, **12**, 4144 (2004)

Quantum Control-Based

Functional Imaging Through Biological Tissue

Experimental Setup




Dela Cruz et al. *PNAS* **101**, 16996 (2004)


2. Progress in Quantum Control

c. Selective Two-Photon Excitation, CARS, SRS, Electronic and Vibrational Control


Data obtained from a scattering surface at 12m standoff distance, 10 μ J/pulse, ~100 μ m neat liquid film ~10mg/cm² Li, Harris, Wrzesinski, Xu, Lozovoy and Dantus, Opt. Express, 16, 592 (2008)




Optical Image



Selective CARS for PS




Selective CARS for **DNT**



“Highly Selective Standoff Detection and Imaging of Trace Chemicals in a Complex Background using Single-Beam Coherent Anti-Stokes Raman Scattering”,
App. Phys. Letters **99**, 101109 (2011)


3. Challenges in Chemistry

a. Chemical Reactions



LASER CHEMISTRY
J. Chem. Educ. 43, 566 (1966)

LASER CHEMISTRY
SCI. AMER. 240, 114 (1979)





Kosloff, Rice, Gaspard, Tersigni, and Tannor propose achieving chemical selectivity by shaping of intense femtosecond pulses, **introduce an iteration procedure for optimization of the pulse shapes** (1989).

Judson and Rabitz propose using a **learning algorithm** to control molecules (1992)

Assion, Baumert, Bergt, Brixner, Kiefer, Seyfried, Strehle, Gerber Control of chemical reactions by feedback-optimized phase shaped laser pulses (1998)

3. Challenges in Coherent Control

a. Chemical Reactions






Pastirk and Dantus JCP 108, 4375 (1998)

Quantum control of a concerted chemical reaction

Dynamic Stark Control of Photochemical Process

Sussman et al. Science 314, 278 (2006)




3. Challenges in Coherent Control



a. Chemical Reactions

iii. Bimolecular reactions


From atoms to molecules ...




Femtosecond Photoassociation



Marvet et al. Chem. Phys. Lett. 245, 393 (1995)



Two-photon photoassociation of hot Mg atoms




Rybak et al. PRL 107, 273001 (2011)

3. Challenges in Coherent Control


a. Chemical Reactions

i. A mol is a very large quantity


ii. Coherent vs. Incoherent



Science **282**, 919 (1998)



Either no coherence or very short lived.



J. Phys. Chem. A **112**, 3789 (2008)



Pastirk et al. *J. Phys. Chem. A* **109**, 2413 (2005)

Dela Cruz et al. *J. Phys. Chem. A* **109**, 8447 (2005)


Ann. Rep. Prog. Chem. C **102**, 227 (2005)

Is photofragmentation pattern phase dependent?

Comprehensive exploration on the role of phase coherence on the fragmentation of large molecules by intense shaped pulses






Zhu et al. J. Phys. Chem. A. 112, 3789 (2008)




3. Challenges in Coherent Control

b. Coherent Control of Electrons, Electronic States and Electronic Transitions



Peak Intensity = 8×10^{13} W/cm²




Peak Intensity = 5×10^{14} W/cm²

3. Challenges in Coherent Control


b. Coherent Control of Electrons, Electronic States and Electronic Transitions

Electronic Wave Packets using $\frac{1}{2}$ -cycle pulses




R. R. Jones, PRL 76, 3927 (1996)

Attosecond electronic wave packet interferometry




Remetter et al, Nature Phys. 2, 323 (2006)



Holler, PRL 106, 123601 (2011)

3. Challenges in Coherent Control

c. Breaking strong chemical bonds while leaving weak ones intact



Kalcic et al. *J. Am. Chem. Soc.* **131**, 940 (2009)

Smith et al. *J Am Soc Mass Spectrom*, **21**, 2031 (2010)


Conclusion: fs-LID is an important method for PTM analysis:

Other methods: Electron capture dissoci., Electron Transfer dissoci., and VUV dissoci.

Perspective: Post-translational modifications: a challenge for proteomics and bioinformatics, Appel, Bairoch, *Proteomics* 6,1525-6 (2004)



Cellular tumor antigen p53: (*Science* Molecule of the year 1993), has more than 18 phosphorylation sites.

DNA winds around Histones, gene regulation and DNA repair depend on histone PTM's.




3. Challenges in Coherent Control

c. Determining the role of coherent quantum processes in nature



Vibrational Coherence in Vision

Wang et al. Science 266, 5184 (1994)






Coherence in Photosynthesis

Engel et al. Nature (2007)

“direct evidence for remarkably long-lived electronic quantum coherence playing an important part in energy transfer processes within this system.”

Coherently wired light-harvesting in photosynthetic marine algae **at ambient temperature**

Collini et al. Nature (2010)



4. Future Outlook and Applications


a. Phase Control of Advanced Sources – towards TL single-cycle pulses

Octave spanning (single-cycle) laser pulses can access any frequency in the spectrum

Typical 100 fs laser, centered at 800 nm

Ultrafast laser producing 12 fs pulses; 2P (x8.3), 3P (x69)

Ultrafast laser producing 4 fs pulses; 2P (x25), 3P (x625)



In 1997 Murnane and Kapteyn reported generation up to the 297th harmonic! =>Coherent X-rays with $\lambda \sim 2$ nm

A 3fs pulse doubles in pulse duration after propagating 12" in air.

Applications of single-cycle pulses require ultra-precise phase measurement and compression

MIIPS has the accuracy and precision to enable this research

4. Future Outlook and Applications

a. Phase Control of Advanced Sources

Femtosecond Free-Electron X-Ray Sources: Sub-10fs pulse duration




Table top x-ray sources:


Compact Coherent X-Ray Sources <10nm wavelength

Table top attosecond laser sources:


Sub-80as (sub-30as) isolated attosecond pulses in the x-ray regime

Ultrafast laser sources:

From single cycle pulses to self-compressing <5fs sources
It is all a matter of phase control.




Single-cycle pulse from Er-doped fiber technology, Krauss et al. (2010)





4. Future Outlook and Applications

a. Phase Control of Advanced Sources – new concepts



Fiber laser oscillators capable of sub-40fs 20nJ/pulse output. Dantus and Wise groups: *Opt. Express* **19**, 12074 (2011)



Sub-30fs Ytterbium oscillator



Synthetic comb sources



Compact <9fs fiber lasers





Filamentation phenomena




A. Braun et al., *Optics Lett.* **20**, 73 (1995)

4. Future Outlook and Applications


a. Phase Control of Advanced Sources – pulse shaping attosecond pulses



CEP phase effect on HHG
Baltuska et al. Nature (2003)



CEP phase effect on HHG
Calegari et al. PRA (2011)




McPherson et al HHG JOSAB (1987); Classical model Corkum PRL (1993),
Popmintchev et al. Nature Photonics Review (2010); Chen et al. PRL (2010)



4. Future Outlook and Applications

b. Quantum control-based molecular sensing and diagnostics

Images of turbulence in a jet of CO_2 in air



Selective Single Beam CARS






J. Raman Spec. **42**, 393-398 (2011)


Collaboration with S. Roy Spectral Energies, and J. Gord (AFRL)

Single-Shot Gas Phase Thermometry via Multi-Pulse Time-to-Frequency Mapping of Coherence Dephasing

Yue et al., *J. Anal. Chem.* (2012)




Spectral Energies, LLC



Femtosecond laser
electronic excitation tagging
R. B. Miles

Appl. Opt. **50**, 5158 (2011)



Lines are written by
the laser and imaged
with a fast gated
camera.







4. Future Outlook and Applications

c. Material Synthesis and Material Processing

Single shot ablation of silicon



In-situ Laser Induced Breakdown Spectroscopy (LIBS)
for real-time monitoring of the machining process





Gunaratne et al. J. App. Phys. 106, 123101(2009)

Gunaratne et al. Chem. Phys. Letts.
423, 197 (2006)

4. Future Outlook and Applications

d. Connecting the macro-micro-and-nano worlds



Spatiotemporal Coherent Control of Lattice Vibrational Waves



Surface-mediated four-wave mixing of nanostructures with counterpropagating surface plasmon polaritons

Liu, Wang, Potma, Opt Lett 36 2348 (2011)

Measurement and Control of Ultrashort Optical Pulse Propagation in Metal Nanoparticle-Covered Dielectric Surfaces



Gunn, Ewald, Dantus, Nano Letters 6, 2804 (2006).


Gunn, High, Lozovoy, Dantus J Phys Chem C 114 12375 (2010)

4. Future Outlook and Applications


e. Coherent control and communications and information processing

Encoding and decoding of femtosecond pulses


Weiner Heritage Salehi, Opt Lett 13 300 (1988)



Asynchronous encrypted information transmission with sub-6 fs laser system at 2.12GHz




Xu, Coello, Nogueira, Cruz, Dantus
Opt Express 16, 15109 (2008)




Experimental coherent
computation of a multiple-
input AND gate using pure
molecular superpositions

Amitay Kosloff Leone Chem
Phys Lett 359 8 (2002)


Optical Code-Division Multiple Access Using Broad-Band Parametrically Generated Light




Peer Dayan Silberberg Friesen J Lightwave
Tech 22 1463 (2004)



From Fundamental Concepts to Applications






Gavin Reid	(MSU, Chemistry, Proteomics)
Daniel Jones	(MSU, Bio Chem, Metabolomics)
Steven P. Arnoczky	(MSU, Lab for Comp. Orthopedic Res.)
George Abela MD	(MSU, Cardiology)
Chong-Yu Ruan	(MSU, ultrafast electron diffraction)
James Resau MD	(Van Andel Institute, cancer)
Stephen Boppart	(UIUC, femtosecond endoscopy)
Jim Gord	(Air Force Research Lab, CARS)
Sukesh Roy	(Spectral Engines LLC, CARS, Machining)
Dominique Fourquette	(Michigan Aerospace, sensing)
Flavio Cruz	(UNICAMP, Brazil, Novel laser sources)
Sunney Xie	(Harvard, CARS microscopy)
Rich Mathis	(Berkeley, low frequency in biomolecules)
Frank Wise	(Cornell, fiber laser design)
Maurice Jansen	(Free Univ. Amsterdam, PEPICO)
Shaul Mukamel	(UC Irvine, NLO Theory)

Phase control enables fundamental research and applications

MICHIGAN STATE
UNIVERSITY




Biophotonic
SOLUTIONS INC.
ULTRAFAST PULSE SHAPING & COMPRESSION

Spectral
Energies, LLC


Backup Slides for Follow Up Discussion

Multiphoton Intrapulse Interference Phase Scan (MIIPS)



Theory



Experiment






5 fs pulse with
500 fs³ TOD





Optics Express **16**, 592 (2008)
JOSA B **25**, A140 (2008)

What is Multiphoton Intrapulse Interference?



Selective nonlinear excitation is useful!



* Albota, M.A., C. Xu, and W.W. Webb. Applied Optics 37,7352 (1998)

4.6 fs laser Bingwei Xu, Yves Coello Optics Express 14, 10939 (2006)


Coello, Xu, and Dantus, Appl. Optics In press (2010)

Pulse shaping for:



Greater signal

Reduced photodamage

Selective excitation



Photoenhancement!






J.Phys.Chem.A 106, (2002)

Optics Express 11, 1695 (2003)

J. Phys. Chem. 108, 53 2004 *Opt. Commun.* 241, 1841 (2008)

PNAS, 101, 16996 (2004)

Experimental Single-Beam CARS Setup



2010 Pulse Shaping Workshop
August 20-22, 2010



Teaching scientists worldwide how to use sub-5fs pulses routinely




2011 Pulse Shaping Workshop @ MSU Aug 19-21. Keynote by Andrew Weiner

High Sensitivity and Selectivity

SUPER-CARS Imaging

Isomer Discrimination




Scientific Goals Achieved: Sensitivity, Selectivity, Imaging

Next step: making it practical


M. Bremer, P. Wrzesinski, N. Butcher, V. Lozovoy and M. Dantus, App. Phys. Letters **99**, 101109 (2011)

Regarding the Reported Selective Synthesis of Toluene from Acetophenone

Original Figure:



Mass spec. obtained for ~100 fs TL pulses



Dashed lines and red peak indicate contaminants



[1] Levis et al., Science 293, 709 (2001).

[2] R. J. Levis and H. A. Rabitz, J. Phys. Chem. A, 106, 6427 (2002)

Comment to J. Phys. Chem. A **113**, 5264 (2009)

Regarding Toluene from Acetophenone Reported in 2004

Original Figure from Graham et al.:



[1] Levis et al., Science 293, 709 (2001).

[2] R. J. Levis and H. A. Rabitz, J. Phys. Chem. A, 106, 6427 (2002)

[3] Graham, Menkir and R. J. Levis, Spectrochimica Acta B 58, 1097 (2004)

Comment to J. Phys. Chem. A **113**, 5264 (2009)

Project from DHS Phase I (cleared)


Determine sensitivity and selectivity in a complex background at standoff distances and evaluate imaging capability.

Looking at the fingerprint region!

Complex background





Samples: Polymer solution with trace amount of DNT spin-coated on substrate.



SUPER-CARS Sensitivity at 1 meter standoff



1s exposure, <4mW average laser power (Raw Data)



**<3 μ g/cm² DNT detection at 1 meter,
in a complex background**

SUPER-CARS Selectivity for standoff imaging

<4mW average laser power



SUPER-CARS Selective excitation ideal for Standoff Imaging Applications



Used by Bartels, *et al* for CARS

Phys. Rev. Lett. **88**, (2002) in multi-beam geometry. GA selective vib. excitation by Bucksbaum.

Spectral imaging by stimulated Raman scattering

Christian Freudiger, Sunney Xie, Nature Photonics (submitted 2010)

Collaboration with Bingwei Xu, and Marcos Dantus



Milestones in Quantum Control

- Young's double-slit experiment (1803), although it is just linear interference

$$P \propto \left| e^{i\phi_1} + e^{i\phi_2} \right|^2 \propto \cos(\phi_1 - \phi_2)$$

- Zewail proposes femtosecond lasers will be able to beat IVR (1980)
- Tannor Kosloff and Rice propose selectivity via control of wave packet evolution (1985-6)

Re Optimization: This result has analogs in electrical engineering and information theory, where it is part of matched filter theory.

Variational optimization of the shape, duration, and separation of the pulses used to generate the selectivity of reactivity, and analysis of the changes induced by inclusion of all degrees of freedom of the molecule (say in the sense of a reaction path Hamiltonian).

- Brumer and Shapiro propose one vs. three-photon quantum control (1986)

$$P \propto \left| e^{i\phi_3} + e^{i3\phi_1} \right|^2 \propto \cos(\phi_3 - 3\phi_1)$$

- Dantus, Rosker and Zewail Femtosecond Transitions State Spectroscopy (1987)
- Kosloff, Rice, Gaspard, Tersigni, and Tannor propose achieving chemical selectivity by shaping of intense femtosecond pulses, and introduce an iteration procedure for optimization of the pulse shapes (1989).
- Bergmann et al. Adiabatic population transfer driven by delayed pulses proposed (1989)

Milestones in Quantum Control

- Chelkowski, Bandrauk Corkum propose and Boers, van Linden van der Heuvell and Noordam demonstrate efficient population transfer in a three-level ladder system by using chirped pulses ChiRAP (1990)
- Judson and Rabitz propose using a learning algorithm to control molecules (1992)
- Bergmann et al. STIRAP experimentally demonstrated (1993)
- Cerullo, Bardeen, Wang and Shank chirped pulse excitation of molecules in solution (1996)
- Gordon Experimental demonstration of one vs. three-photon control (1997)
- Weinacht, Ahn and Bucksbaum “read-write” on sculpted Rydberg wave packet (1998)
- Pastirk, Brown, Zhang and Dantus chirp control of a chemical reaction (1998)
- Silberberg demonstrates multi-chromatic two photon control (1998)






$$P(2\omega) \propto \left| 1 + e^{i\phi(\omega-\Delta)+i\phi(\omega+\Delta)} \right|^2$$

- GA assisted pulse compression (1997)
- Assion, Baumert, Bergt, Brixner, Kiefer, Seyfried, Strehle, Gerber Control of chemical reactions by feedback-optimized phase shaped laser pulses (1998)

Input field Parameters Load from file(s)

Time Domain

SHG Stim. Raman MIIPS MIIPS II FROG/XFROG AC



Phase Control

$$\varphi(\omega) = \varphi_0 + \varphi_1 \cdot (\omega - \omega_0) + \frac{\varphi_2}{2!} \cdot (\omega - \omega_0)^2 + \dots$$




Polynomial Cosine/Sine Binary

Amplitude Control

Function:
activated pixel - 1,
otherwise - 0.



Binary Cos^2 Cos^4








About Version Activation

Input field Parameters Load from file(s)


Time Domain

SHG Stim. Raman MIIPS MIIPS II FROG/XFROG AC



Phase Control

$$\varphi(\omega) = \varphi_0 + \varphi_1 \cdot (\omega - \omega_0) + \frac{\varphi_2}{2!} \cdot (\omega - \omega_0)^2 + \dots$$



Polynomial

Cosine/Sine

Binary

Amplitude Control

Function:
activated pixel - 1,
otherwise - 0.



Binary

Cos^2






Cos^4

BSI - Copyright © 2005-2010

Input field Parameters Load from file(s)

Time Domain

SHG Stim. Raman MIIPS MIIPS II FROG/XFROG AC



Phase Control

$$\varphi(\omega) = \varphi_0 + \varphi_1 \cdot (\omega - \omega_0) + \frac{\varphi_2}{2!} \cdot (\omega - \omega_0)^2 + \dots$$



Polynomial Cosine/Sine Binary


Amplitude Control

Function:
activated pixel - 1,
otherwise - 0.





Binary Cos^2 Cos^4







Sample of random 16-bit binary phases




Is molecular response predictable?




Effect of phase & amplitude shaping




Generalizing to other molecules





Back to MS
fragmentation in
strong fields





Xin Zhu

Either no e-coherence
or very short lived.


Searching for coherence

Coherence found on acetophenone



Zhu, X., Lozovoy, V. V., Shah, J. D. & Dantus, M. "Photodissociation Dynamics of Acetophenone and Its Derivatives with Intense Nonresonant Femtosecond Pulses." *J. Phys. Chem. A* **115**, 1305-1312 (2011).


Part III. 6. The need for pulse compression



Pulse Compression

The goal is to take a broad-bandwidth pulse with highly nonlinear dispersion (quadratic, cubic, etc...) and to compensate the phase distortions to recover a TL pulse.




Pulse after single bounce from dielectric coated mirror



D. Pestov, V. V. Lozovoy, and M. Dantus "Single-beam shaper-based pulse characterization and compression using MIIPS sonogram", Opt. Letters **35**, 1422-1424 (2010)

GA. Pulse Compression

Works but it is slow and not very accurate because search space near TL is flat



Optics Letters **22**, 1793 (1997)

Appl. Phys. B. **65**, 779 (1997)

About evolutionary learning algorithms

The search space = $(P \times A)^N$

N = # of pixels

P = # of phases

A = # Amplitudes

For:

N = 100

P = 100

A = 10

Search space = 10^{300}

10^{200} for phase only

Limited application in the field!

- Size of parameter space vs. sampling (10^8 per day, 10^{12} per life)
- Low repeatability and portability of results
- The search space: Is it convex or needle in the haystack?

Conclusion:

We need a systematic approach

